

FOREST RESEARCH LABORATORY

RESEARCH NOTE 65

a simple model of host resistance to bark beetles

Richard H. Waring
Gary B. Pitman

While studying why bark beetles attack certain pine trees and ignore others, we hypothesized that the amount of carbohydrate reserves available basically controls host resistance. Apparently the monoterpene chemistry of the phloem which attracts beetles differs when reserves are high or low. Moreover, the available carbohydrate reserves also determine the ability of a tree to produce oleoresin and other chemicals which protect it against beetles and subsequent blue stain.

A tree has certain requirements that must be satisfied if it is to live. The roots (Harris et al. 1978, Fogel and Hunt 1979), shoots (Gordon and Larson 1968, Rangnekar and Forward 1973), and even cones (Eis et al. 1965) take priority over stemwood growth. Thus, a decline in stemwood production should indicate a limitation by carbohydrate reserves—and, therefore, increased susceptibility to bark beetles.

Because large trees usually grow more stemwood each year than smaller ones, we faced the

problem of generalizing the relative extent of carbohydrate reserves available. Growth in stemwood volume or biomass can accurately be estimated by extracting a small core from the stem and determining the cross-sectional area grown in the current year. Because foliage biomass or leaf area correlates linearly with the cross-sectional area of conducting tissue at breast height or 1.37 m (Grier and Waring 1974), we normalized growth in terms of a square meter (m^2) of foliage. We thus proposed that trees with high resistance would produce a relatively large amount of sapwood each year, whereas susceptible trees would produce much less. In lodgepole pine (*Pinus contorta* Dougl. ex Loud.), we can directly convert this ratio of basal-area growth/sapwood basal-area to stemwood biomass (kg/m^2) of foliage by multiplying by 0.87 (Gholz et al. 1979).

We wished to establish the number of beetle attacks required to kill trees at different levels of resistance—that is, having different ratios of basal-area growth/sapwood basal-area. After beetles

attacked plots near our study in 1978, we surveyed 75 trees by measuring the diameters at breast height, counting the attacks in a 30-cm band (± 15 cm from breast height) encircling the tree, then calculating the beetle attacks per square meter of bark surface.

Figure 1 summarizes our findings. None of the attacked trees had vigor indices above 15 percent (15% of the sapwood laid down in the current year). No trees with vigor indices above 8 percent were killed because attack densities did not exceed $200/m^2$. Trees with vigor indices below 8 percent could be killed if beetles attacked in densities equal to or greater than about 24 times the vigor index. For example, trees with vigor indices of 5 percent could be killed by attacks exceeding 120 beetles/ m^2 .

Having modified the vigor index of hundreds of trees by thinning, fertilizing, and stressing with additions of sugar and sawdust, we now are testing the accuracy of the model to predict mortality on more than 100 recently attacked trees.




Figure 1.

Mortality of lodgepole pine in relation to intensity of beetle attack and tree vigor index.

literature cited

EIS, S., E.H. GARMAN, and L.F. EBELL. 1965. Relation between cone production and diameter increment of Douglas-fir [*Pseudotsuga menziesii* (Mirb.) Franco], grand fir [*Abies grandis* (Dougl.) Lindl.], and western white pine (*Pinus monticola* Dougl.). Canadian Journal of Botany 43:1553-1559.

FOGEL, R., and G. HUNT. 1979. Fungal and arboreal biomass in a western Oregon Douglas-fir ecosystem: distribution patterns and turnover. Canadian Journal of Forest Research 9:245-256.

GHOLZ, H.L., C.C. GRIER, A.G. CAMPBELL, and A.T. BROWN. 1979. Equations for estimating biomass and leaf area of plants in the Pacific Northwest. Oregon State University, Forest Research Laboratory, Corvallis. Research Paper 41. 37 p.

GORDON, J.C., and P.R. LARSON. 1968. Seasonal course of photosynthesis, respiration, and distribution of ^{14}C in young *Pinus resinosa* trees as related to wood formation. Plant Physiology 43:1617-1624.

GRIER, C.C., and R.H. WARING. 1974. Conifer foliage mass related to sapwood area. Forest Science 20:205-206.

HARRIS, W.F., R.S. KINERSON, and N.T. EDWARDS. 1978. Comparison of belowground biomass of natural deciduous forest and loblolly pine plantations. Pedobiologia 17:369-381.

RANGNEKAR, P.V., and D.F. FORWARD. 1973. Foliar nutrition and wood growth in red pine: effects of darkening and defoliation on the distribution of ^{14}C -photosynthate in young trees. Canadian Journal of Botany 51:103-108.

the authors

R. H. Waring is a professor and G. B. Pitman is an associate professor, Department of Forest Science, School of Forestry, Oregon State University, Corvallis, Oregon 97331.

acknowledgments

Support for this research came from McIntyre-Stennis funds and a National Science Foundation Grant (DEB-7904540). We thank Steve Hyland and Lance Waring for help in the field.

forest research laboratory
school of forestry
oregon state university
corvallis, oregon 97331

Non-Profit Org.
U.S. Postage
PAID
Permit No. 200
Corvallis, OR 97331

As an affirmative action institution that complies with Section 504 of the Rehabilitation Act of 1973, Oregon State University supports equal educational and employment opportunity without regard to age, sex, race, creed, national origin, handicap, marital status, or religion.